

WARNING


READ LABEL BEFORE USE. KEEP OUT OF REACH OF CHILDREN. FOR ANIMAL TREATMENT ONLY.

20L

Inovata[®] **TISON ATTACK**

POUR ON/SPOT ON

For the treatment and control in crossbred sheep of susceptible gastrointestinal roundworms and lungworm, including strains with single resistance to macrocyclic lactones and levamisole.

2**Abamectin
Levamisole****MADE
HERE.
FOR
HERE.****ACTIVE INGREDIENTS:**

15 g/L abamectin and 26 g/L levamisole

Inovata[®]

TISON ATTACK Inovata®

Read label and obtain special instructions before use. Do not handle until all safety precautions have been read and understood.

WARNING
Harmful if swallowed. Causes serious eye irritation. May cause an allergic skin reaction. Suspected of causing genetic defects. May cause damage to organs. Suspected of damaging fertility or the unborn child. May cause harm to breast-fed children.

Handling Precautions: Wear protective gloves/ clothing, eye/ face protection. Wash hands thoroughly after handling. Contaminated work clothing should not be allowed out of the workplace. Use personal protective equipment as required. Do not breathe fume/vapours. Avoid contact during pregnancy/ while nursing. Do not eat, drink or smoke when using this product.

First Aid: If medical advice is needed, have product container or label at hand. IF SWALLOWED: Call a POISON CENTER or doctor/ physician if you feel unwell. Rinse mouth. IF ON SKIN: Wash with plenty of soap and water. If skin irritation or rash occurs: Get medical advice/attention. IF IN EYES: Rinse cautiously with water for several minutes. Remove contact lenses if present and easy to do so. Continue rinsing. If eye irritation persists: Get medical advice/attention. Wash contaminated clothing before reuse. If exposed, concerned or feel unwell: Call a POISON CENTER or doctor/physician.

National Poison Centre 24hr Phone: 0800 POISON (0800 764 766)

EFFICACY SPECTRUM:
Tison Attack contains two broad-spectrum anthelmintics from different drench families. Abamectin is a member of the macrocyclic lactone drench family and levamisole is an imidazothiazole. Tison Attack is effective against strains of internal parasites sensitive to one or both of the anthelmintics in this drench:

Gastrointestinal roundworms:

- Barber's Pole Worm
- Haemonchus contortus**
- Small Brown Stomach Worm
- Teladorsagia circumcincta**
- Stomach Hair Worm
- Trichostrongylus axei*
- Black Scour Worms
- Trichostrongylus vitrinus*,
- T. colubriformis*
- Thread Worm
- Strongyloides papillosus*
- Small Intestinal Worms
- Cooperia curticei**
- Thin-Necked Intestinal Worms
- Nematodirus filicollis*,
- N. spathiger*
- Large Bowel Worm
- Oesophagostomum venulosum*
- Large-mouthed Bowel Worm
- Chabertia ovina*
- Whipworm
- Trichuris ovis*
- Lungworm:**
- Dictyocaulus filaria*
- *includes inhibited fourth stage larvae.

NOTE: Resistance can develop to any anthelmintic. Ask your local veterinary practitioner or animal health adviser for recommended parasite management practices for your local area to reduce the potential for resistance to develop. It is advisable that a resistance test be conducted regularly when using any parasite treatment.

DIRECTIONS FOR USE:
Dosage and Administration:
By law, the user must take due care, seeking expert advice if necessary, to avoid unnecessary pain or distress in the treated animal when using the product in a manner other than as recommended on the label.

For use in a strategic worm control program: speak to your animal health adviser or veterinarian. Correct drenching technique should be used.

For topical application only. ADMINISTER DIRECTLY ONTO THE SKIN ON THE BACKLINE. This product should be applied

on clean and healthy skin. Do not apply to animals with skin lesions or damaged skin. This product should not be used at the same time as topical fly or lice control products. **Off-shears and within 2 weeks of shearing** apply with a drench gun nozzle as a single 200-300mm midline stripe down the backbone in the middle of the back.

If greater than 2 weeks after shearing divide the dose approximately in half and apply it as two individual spots. The spots should be 100-150 mm apart, along the backbone and centred in the middle of the back. A specially designed nozzle can be used to do this with one ejection of the drench gun (see picture).

IMPORTANTLY, THE DRENCH GUN NOZZLE MUST BE PUSHED THROUGH FLEECE TO THE SKIN LEVEL

A representative sample of animals should be weighed before treatment. Dose the mob to the heaviest animal by liveweight in each group (ewes, wethers, rams, lambs). Do not underdose. If there is a large variation in size within the group, draft into two or more lines based on bodyweight to avoid excessive overdosing. DO NOT exceed the stated dose, or dose more frequently than 3-weekly. Check accuracy of drenching equipment regularly. Clean gun thoroughly before and after use.

Dose Volume Table:

1 mL per 2.5 kg live-weight (6mg/kg abamectin and 10mg/kg levamisole)
The effect of rainfall on the efficacy of Tison Attack has not been assessed. It is recommended that it is not applied on wet sheep or if rain is imminent. **Do not use in lambs under 6 weeks of age and under 16 kg liveweight**

Environmental Protection: Very toxic to aquatic life. Harmful to the soil organisms. Hazardous to terrestrial vertebrates. Hazardous to terrestrial invertebrates. Avoid release to the environment. Collect spillage.

Disposal: Preferably dispose of product by use. Otherwise dispose of product and packaging at an approved landfill facility. Triple rinse container. Recycle empty rinsed container at an approved recycling centre. Otherwise puncture or crush empty container and dispose of at approved landfill facility. Do not use container for any other purpose. For details on how to correctly prepare your container and collection sites please visit the Agrecovery website www.agrecovery.co.nz

Emergency Response: In case of spillage, wear appropriate protective clothing and prevent material from entering water ways. Absorb spills with inert material and place in waste container. Sweep up solids without creating dust. Wash area with water and absorb with further inert material. Dispose of waste safely.

DRUM TAP ENCLOSED
Fit to internal thread after removing small drum cap.

TWIN APPLICATOR NOZZLE ENCLOSED



Cut along dotted line

DOSE VOLUME TABLE:

Live-weight (kg)	Dose (mL)	20L treats (head)
16 - 20	8	2500
21 - 25	10	2000
26 – 30	12	1666
31 - 35	14	1428
36 – 40	16	1250
41 – 45	18	1111
46 – 50	20	1000
51 - 55	22	909
56 - 60	24	833
Sheep in excess of 60 kg must be dosed at 1mL per 2.5 kg liveweight.		

DOSE RATE 1 mL per 2.5 kg liveweight

TISON ATTACK



ENVIRONMENTALLY HAZARDOUS SUBSTANCE,
LIQUID N.O.S.
(Contains Abamectin and Levamisole)
UN No. 3082, Packaging Group III, HazChem 3Z



Inovata®

Registered to:
Inovata Ltd
16 Sheffield Crescent, Burnside, Christchurch 8053
PO Box 20449, Bishopdale, Christchurch 8543
www.inovata.co.nz
sales@inovata.co.nz
Freephone 0800 942 006

Batch:
Expiry: