

WARNING - FLAMMABLE LIQUID

READ LABEL BEFORE USE. KEEP OUT OF REACH OF CHILDREN. FOR ANIMAL TREATMENT ONLY.

20L

Inovata[®]

Abamectin

POUR-ON FOR CATTLE

For the treatment and control of internal and external parasites of cattle.



1

Abamectin

**MADE
HERE.
FOR
HERE.**



ACTIVE INGREDIENTS:

Abamectin 10 mg/ml. Contains >941 g/L Diethylene glycol monobutyl ether.

Inovata[®]

Abamectin

POUR-ON

Inovata®

WARNING

READ ENTIRE LABEL BEFORE USE

• **Dangerous to the Environment**

HANDLING PRECAUTIONS

Keep away from heat/sparks/open flames/hot surfaces. Wear protective gloves/protective clothing/eye protection. Keep closed when not in use. May be harmful if swallowed, inhaled or absorbed through the skin. May cause mild skin and eye irritation. Repeated exposure may cause skin allergy. Abamectin may possibly affect development and/or reproduction and may have effects on/or via lactation. Diethylene glycol monobutyl ether may possibly affect kidney if swallowed and liver if inhaled. Wash hands and exposed skin before meals and after use. Do not eat, Drink

or smoke when using this product. Do not inhale vapor. Avoid skin and eye contact. Very toxic to aquatic organisms and terrestrial invertebrates. Toxic to the soil environment. Harmful to terrestrial vertebrates. Avoid contamination of any water supply with product or empty container. Avoid release to the environment.

FIRST AID If swallowed do NOT induce vomiting. If skin or hair contact occurs remove contaminated clothing and flush skin and hair with running water. If splashed in eyes wash out immediately with water. For advice contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately.

EMERGENCY RESPONSE In case of spillage wear appropriate protective clothing. Prevent material from entering waterways and absorb spills with inert material and place in waste container. Wash area and absorb with further inert material.



ENVIRONMENTALLY HAZARDOUS SUBSTANCE,
LIQUID, N.O.S. (Contains Abamectin)
UN No. 3082, Packaging Group III, HazChem 3Z

INDICATION Inovata Abamectin Pour-On for Cattle, when used at the recommended dose level of 500 micrograms Abamectin per kg liveweight, effectively controls the following parasites of cattle:

- Gastro-intestinal roundworms: Adult and immature *Ostertagia spp* (including inhibited immatures), *Haemonchus spp*, *Trichostrongylus spp*, *Cooperia spp*, *Bunostomum spp*, *Oesophagostomum spp*, and adults of *Chabertia spp*, *Nematodirus spp*, *Trichuris spp*, and *Strongyloides spp*.
- Lungworms: Adult and immature *Dictyocaulus viviparus*.
- Biting and Sucking Lice: *Bovicola bovis*, *Linognathus vituli*.

DIRECTIONS FOR USE By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

Inovata Abamectin Pour-On for Cattle contains 10 mg/mL abamectin, a member of the macrocyclic lactone group of compounds. It is effective against internal and external parasites sensitive to this family. Resistance can develop to any anthelmintic. Check the effectiveness of anthelmintics at least annually. Follow recommended parasite management practices to reduce the potential for resistance to develop. Check accuracy of pour-on applicator regularly. A representative sample of animals should be weighed before treatment. Dose rate to be based on heaviest cattle in each group (bulls, cows, steers, calves etc.). Do not under-dose. Where there is a large variation in size within the group, draft into two or more lines based on bodyweight, to avoid excessive overdosing. Ready to use with standard pour-on equipment. Apply along the backline from shoulders to tail. Inovata Abamectin Pour-On is rainfast, even if rain falls 2 hours or more after treatment.

DOSAGE It is strongly recommended not to use this product in cattle under four months of age and/or 100 kg live weight. The dose rate for cattle is 500 µg Abamectin per kg bodyweight, which is provided by 1 mL Inovata Abamectin Pour-On per 20 kg bodyweight, or according to the following table:

Weight Range	Dose Volume	Dose per 20 L Pack	Weight Range	Dose Volume	Dose per 20 L Pack
			351-400 kg	20 mL	1000
101-150 kg	7.5 mL	2666	401-450 kg	22.5 mL	888
151-200 kg	10 mL	2000	451-500 kg	25 mL	800
201-250 kg	12.5 mL	1600	501-550 kg	27.5 mL	727
251-300 kg	15 mL	1333	551-600 kg	30 mL	666
301-350 kg	17.5 mL	1142	601-650 kg	32.5 mL	615

Heavier animals (e.g. mature bulls) should receive an additional 1 mL for each 20 kg over 650 kg.

DRUM TAP ENCLOSED.

Fit to internal thread after removing small drum cap.



Cut along dotted line.

WITHHOLDING PERIODS It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

MILK: Milk intended for sale for human consumption must be discarded during treatment and for not less than **35 days** following the last treatment.

MEAT: Cattle producing meat and offal for human consumption must not be sold for slaughter during or within **35 days** of the last treatment.

STORAGE Store below 25°C. Protect from sunlight and keep it well closed.

DISPOSAL Preferably dispose of product by use. Otherwise dispose of product, packaging and waste at an approved landfill or equivalent facility. Triple rinse empty container, crush or puncture and bury in a suitable landfill, or if appropriate recycle.



Registered pursuant to the ACM Act 1997, No A011958
See www.foodsafety.govt.nz for registration conditions.

Approved pursuant to the HSNO Act 1996, approval code HSR100682.
See www.epa.govt.nz for controls.

Registered to:

Inovata Ltd, 16 Sheffield Crescent, Burnside, Christchurch 8053
Freephone: 0800 942 006
www.inovata.co.nz

Batch:
Expiry: