

**DANGER**

READ LABEL BEFORE USE. KEEP OUT OF REACH OF CHILDREN. FOR ANIMAL TREATMENT ONLY.

**10L**


# EraserTape

## TRIPLE ACTIVE COMBINATION PLUS TAPE ORAL DRENCH FOR SHEEP

For the treatment and control of mature stages of all major gastrointestinal parasites in sheep, including those with single or dual resistance to benzimidazole, levamisole/morantel and avermectin/milbemycin families. Also treats and controls lungworm, itch mite, nasal bot, tapeworm, and as an aid in the control of adult liver fluke infestation in sheep.

**3**

Abamectin  
Levamisole Hydrochloride  
Albendazole



**ACTIVE INGREDIENTS:** Abamectin 1 g/L; Levamisole Hydrochloride 40 g/L; Albendazole 25 g/L; Praziquantel 19 g/L; Copper 2.1 g/L; **Selenium** 0.5 g/L; Zinc 0.6 g/L; Cobalt 0.21 g/L and Iodine 0.76 g/L.

**Inovata**<sup>®</sup>

**DANGER**

READ LABEL BEFORE USE. KEEP OUT OF REACH OF CHILDREN. FOR ANIMAL TREATMENT ONLY.

**10L**

# EraserTape

## TRIPLE ACTIVE COMBINATION PLUS TAPE ORAL DRENCH FOR SHEEP

For the treatment and control of mature stages of all major gastrointestinal parasites in sheep, including those with single or dual resistance to benzimidazole, levamisole/morantel and avermectin/milbemycin families. Also treats and controls lungworm, itch mite, nasal bot, tapeworm, and as an aid in the control of adult liver fluke infestation in sheep.

**3**

Abamectin  
Levamisole Hydrochloride  
Albendazole



**ACTIVE INGREDIENTS:** Abamectin 1 g/L; Levamisole Hydrochloride 40 g/L; Albendazole 25 g/L; Praziquantel 19 g/L; Copper 2.1 g/L; **Selenium** 0.5 g/L; Zinc 0.6 g/L; Cobalt 0.21 g/L and Iodine 0.76 g/L.

**Inovata**<sup>®</sup>

### READ ENTIRE LABEL BEFORE USE

Eraser Tape contains albendazole, a member of the benzimidazole family, levamisole, a member of the imidazothiazole family and abamectin, a member of the macrocyclic lactone family. It is effective against gastrointestinal worms sensitive to one, two or all three broad spectrum drench families. It also contains praziquantel, a member of the isoquinoline family which is effective against tapeworm. Provides a supplemental source of copper, **selenium**, zinc, iodine and cobalt.

**INDICATIONS** Resistance may develop to any chemical. It is advisable that a resistance test be conducted regularly when using any parasite treatment. Ask your local veterinary practitioner or animal health adviser for recommended parasite management practices for your area to reduce development of resistance.

**Sheep:** For the effective treatment and control of mature and immature stages of the following gastrointestinal parasites (\*includes inhibited fourth stage larvae):

*Teladorsagia circumcincta\**, *Teladorsagia trifurcata\**, *Haemonchus contortus\**, *Trichostrongylus axei*, *Trichostrongylus colubriformis*, *Trichostrongylus vitrinus*, *Cooperia curticei\**, *Cooperia oncophora\**, *Cooperia punctata\**, *Trichuris ovis*, *Chabertia ovina*, *Oesophagostomum venulosum*, *Strongyloides papillosus*, *Nematodirus filicollis* and *Nematodirus spathiger*.

Also effective against all stages of tapeworm (*Moniezia expansa*), lungworm (*Dictyocaulus filaria*), itch mite (*Psorobia ovis*), nodule worm (*Oesophagostomum columbianum*) and all larval stages of nasal bot (*Oestrus ovis*) and as an aid in the control of adult liver fluke infestations in sheep.

**Anthelmintic resistant strains:** For the effective treatment and control of those gastrointestinal worms that have single and dual resistance to this combination of benzimidazole, levamisole/ morantel or avermectin/milbemycin anthelmintic action families.

**DIRECTIONS FOR USE** By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

**SHAKE WELL BEFORE USE.** For oral administration only. Ensure formulation remains evenly suspended during use. A representative sample of animals should be weighed before treatment. Draft animals into lines of similar bodyweight, then dose according to the weight of the heaviest animal in the line. When there is a large variation in size within the group, dose rates should be based on the label directions for each weight range. Drafting into two or more lines may be appropriate to avoid excessive overdosing. Do not underdose. Correct drenching technique should be used. Check accuracy of drench guns regularly. Rinse gun with water immediately after use. Accuracy of drench guns should be checked before and during drenching. Ensure that the dosing technique is accurate and that each sheep receives and swallows the full dose.

### DOSE RATE

1 ml per 5 kg bodyweight equivalent to albendazole 5 mg, levamisole hydrochloride 8 mg, abamectin 0.2 mg, praziquantel 3.8 mg, **selenium** 0.1 mg per kg bodyweight.

Bodyweight:	Dose	Doses per 10 L pack
15 kg	3 mL	3333
16-20 kg	4 mL	2500
21-25 kg	5 mL	2000
26-30 kg	6 mL	1667
31-35 kg	7 mL	1429
36-40 kg	8 mL	1250
41-45 kg	9 mL	1111
46-50 kg	10 mL	1000
51-55 kg	11 mL	909
56-60 kg	12 mL	833
61-65 kg	13 mL	769
66-70 kg	14 mL	714
71-75 kg	15 mL	667
Over 75 kg add 1 mL for every 5 kg bodyweight		

### WITHHOLDING PERIODS

**It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.**

Sheep producing meat or offal for human consumption must not be sold for slaughter either during treatment or within 21 days of the last treatment;

Milk intended for human consumption must be discarded during treatment and for not less than 35 days following last treatment.

### STORAGE

Store locked up below 25°C, out of direct sunlight.

### DISPOSAL

Preferably dispose of product by use. Otherwise dispose of product and packaging at an approved landfill facility. Triple rinse container. Recycle empty rinsed container at an approved recycling centre. Otherwise puncture or crush empty container and dispose of at approved landfill facility. Do not use container for any other purpose.



**Registered to:** Inovata Ltd  
16 Sheffield Crescent, Burnside, Christchurch 8053  
PO Box 20449, Bishopdale, Christchurch 8543  
www.inovata.co.nz | sales@inovata.co.nz  
Freephone 0800 942 006

EPA Approval number HSR100758  
See www.epa.govt.nz for approval conditions.  
Registered pursuant to the ACVM Act 1997: A011948  
See www.foodsafety.govt.nz for registration conditions.

Harmful if swallowed. Harmful in contact with skin. Harmful if inhaled. May cause allergic skin reactions. Suspected of causing genetic defects. May damage fertility or the unborn child. May cause harm to breast-fed children. May cause damage to organs.

**HANDLING PRECAUTIONS** Avoid breathing mist/vapours/spray. Do not eat, drink or smoke when using this product. Contaminated clothing should not be allowed out of the workplace. Wear protective gloves/clothing. Use personal protective equipment as required. Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Avoid contact during pregnancy/while nursing. Wash hands thoroughly after handling. Avoid release to the environment.

**FIRST AID** If medical advice needed, have container or label at hand. IF SWALLOWED: Call a POISON CENTER (0800 764 766) or doctor/physician if you feel unwell. Rinse mouth. IF ON SKIN: Wash with plenty of soap and water. If skin irritation or rash occurs get medical advice/attention. Wash contaminated clothing before reuse. IF exposed or concerned: Get medical advice/attention.

**ENVIRONMENTAL PROTECTION** Very toxic to aquatic life with long lasting effects. Harmful to terrestrial vertebrates. Toxic to terrestrial invertebrates.

### WARNINGS AND CONTRAINDICATIONS

- Do not use on animals which are severely stressed or in ill health
- Do not exceed stated dose or frequency without consulting veterinarian
- Do not use in lambs under 15 kg or under 4 weeks of age
- Doses of 3 or more times those recommended can cause symptoms of levamisole toxicity, so estimate live weights carefully. Dehydrated animals may be more susceptible to toxicity
- Do not treat sheep at intervals of less than 3 weeks, due to **selenium** content
- Do not use at the same time as any other **selenised** fertilizer, prill or product and do not exceed the stated dose or frequency without consulting a veterinarian
- In sheep, liver levels of copper may be quite variable. Consequently there is always a risk of copper poisoning (deaths) occurring following copper supplementation in this species
- Fatal interactions may occur between levamisole and organophosphate dips.

# Inovata®



ENVIRONMENTALLY HAZARDOUS SUBSTANCE  
(Contains Abamectin, **Selenium**)



NEW CODE TO COME 0000-000-000