

**DANGER**

READ LABEL BEFORE USE. KEEP OUT OF REACH OF CHILDREN. FOR ANIMAL TREATMENT ONLY.

**10L**


# Inovata<sup>®</sup> Tape

## DOUBLE ACTIVE PLUS TAPE ORAL DRENCH FOR SHEEP

For the treatment and control in sheep of all major gastrointestinal parasites, including those with resistance to the benzimidazole or levamisole/morantel families. Also treats and controls lungworms and tapeworms, and is an aid in the control of adult fluke. Provides a supplemental source of copper, **selenium**, zinc, iodine and cobalt.

**2**

Levamisole Hydrochloride  
Albendazole



**Inovata<sup>®</sup>**

**ACTIVE INGREDIENTS:** Levamisole Hydrochloride 40 g/L, Albendazole 25 g/L, Praziquantel 19 g/L, Copper 2.1 g/L, **Selenium** 0.5 g/L, Zinc 0.6 g/L, Cobalt 0.21 g/L and Iodine 0.76 g/L.

**DANGER**

READ LABEL BEFORE USE. KEEP OUT OF REACH OF CHILDREN. FOR ANIMAL TREATMENT ONLY.

**10L**


# Inovata<sup>®</sup> Tape

## DOUBLE ACTIVE PLUS TAPE ORAL DRENCH FOR SHEEP

For the treatment and control in sheep of all major gastrointestinal parasites, including those with resistance to the benzimidazole or levamisole/morantel families. Also treats and controls lungworms and tapeworms, and is an aid in the control of adult fluke. Provides a supplemental source of copper, **selenium**, zinc, iodine and cobalt.

**2**

Levamisole Hydrochloride  
Albendazole



**Inovata<sup>®</sup>**

**ACTIVE INGREDIENTS:** Levamisole Hydrochloride 40 g/L, Albendazole 25 g/L, Praziquantel 19 g/L, Copper 2.1 g/L, **Selenium** 0.5 g/L, Zinc 0.6 g/L, Cobalt 0.21 g/L and Iodine 0.76 g/L.

**EFFICACY SPECTRUM** Inovata Tape is intended for use against benzimidazole and levamisole sensitive internal parasites of the following species: *Ostertagia circumcincta*, *Haemonchus contortus*, *Trichostrongylus axei*, *Trichostrongylus colubriformis*, *Cooperia curticei*, *Trichuris ovis*, *Chabertia ovina*, *Oesophagostomum venulosum*, *Strongyloides papillosus*, *Nematodirus filicollis*, *Nematodirus spathiger*, *Bunostomum trigonocephalum*. LUNGWORMS: Large lungworm (*Dictyocaulus filarial*). TAPEWORMS: Heads and Segments (*Moniezia spp.*). AS AN AID IN CONTROL OF LIVER FLUKE (Adults): *Fasciola hepatica*. OVICIDAL: Reduces the output of viable worm and fluke eggs.

**NOTE:** Resistance can develop to any Anthelmintic. Ask your local veterinary practitioner or animal health adviser for recommended parasite management practices for your local area to reduce the potential for resistance to develop. It is advisable that a resistance test be conducted regularly when using any parasite treatment.

**DIRECTIONS FOR USE** By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label. For use in a strategic worm control program: speak to your animal health adviser or veterinarian.

**SHAKE CONTAINER WELL BEFORE USE.** Administer orally and check accuracy of drenching equipment regularly. Clean gun thoroughly before and after use. A representative sample of animals should be weighed before treatment. Dose the mob to the heaviest animal by live-weight in each group (ewes, wethers, rams, lambs). Do not underdose. Where there is a large variation in size within the group, dose rate should be based on the label directions for each weight range. If there is a large variation in size within the group, draft into two or more lines based on bodyweight to avoid excessive overdosing. Correct drenching technique should be used.

**DO NOT** use at the same time as any other **selenised** fertiliser, prill or product without consulting a veterinarian. DO NOT exceed the stated dose, or dose more frequently than 3-weekly.

**CAUTION:** In sheep, liver levels of copper may be quite variable. Consequently, there is always a risk of copper poisoning (deaths) occurring following copper supplementation in this species. Doses of 3 or more times those recommended can cause symptoms of Levamisole toxicity, so estimate live-weights carefully. Dehydrated animals may be more susceptible to toxicity. Fatal interactions may occur between Levamisole and Organophosphate dips.

**DOSE RATE**  
1 mL per 5 kg live-weight

Live-weight (kg)	Dose (mL)	10 L Treats (head)
15 kg	3 mL	3333
16-20 kg	4 mL	2500
21-25 kg	5 mL	2000
26-30 kg	6 mL	1666
31-35 kg	7 mL	1428
36-40 kg	8 mL	1250
41-45 kg	9 mL	1111
46-50 kg	10 mL	1000
51-55 kg	11 mL	909
56-60 kg	12 mL	833
61-65 kg	13 mL	769
66-70 kg	14 mL	714
71-75 kg	15 mL	666

1 mL per 5 kg bodyweight equivalent to levamisole hydrochloride 8 mg, albendazole 5 mg, praziquantel 3.8 mg and selenium 0.1 mg per kg bodyweight.

**WITHHOLDING PERIODS**

It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

**MEAT:** Animals producing meat or offal for human consumption must not be sold for slaughter either during treatment or within 14 days of cessation of the last treatment.

**MILK:** Milk intended for safe human consumption or manufacture must be discarded during treatment and for 35 days following the last treatment.

**STORAGE**

Store below 25°C in original container tightly closed and in a locked, dry and well-ventilated area, away from foodstuffs. Do not freeze.

EPA Approved number HSR100759

See [www.epa.govt.nz](http://www.epa.govt.nz) for approval conditions.

Registered pursuant to the ACVM Act 1997, No A011965

See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions.

ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S.  
(Contains: Albendazole, Sodium Selenate)



Read label and obtain special instructions before use. Do not handle until all safety precautions have been read and understood.

**HANDLING PRECAUTIONS** May be harmful if swallowed. May cause allergy or asthma symptoms or breathing difficulties if inhaled. May cause an allergic skin reaction. Suspected of causing genetic defects. Suspected of causing cancer. May damage fertility or the unborn child. May cause damage to organs. Do not breathe dust/fumes/gas/mist/spray. In case of inadequate ventilation wear respiratory protection. Contaminated work clothing should not be allowed out of the workplace. Wear protective gloves. Use personal protective equipment as required.

**FIRST AID** If medical advice is needed, have product container or label at hand. Call a POISON CENTER or doctor/physician if you feel unwell. **IF INHALED:** If breathing is difficult, remove to fresh air and keep at rest in a position comfortable for breathing. If experiencing respiratory symptoms: Call a POISON CENTER or doctor/physician. **IF ON SKIN:** Wash with plenty of soap and water. If skin irritation or rash occurs: Get medical advice/attention. Wash contaminated clothing before reuse. If exposed, concerned or feel unwell: Get medical advice/attention.

**ENVIRONMENTAL PROTECTION** Harmful to aquatic life with long lasting effects. Harmful to soil environment. Avoid release to the environment.

**DISPOSAL** Preferably dispose of product by use. Otherwise treat the product so that it is no longer toxic and dispose of product and packaging at an approved landfill or other approved facility. Triple rinse empty container, crush or puncture and bury in a suitable landfill, or recycle if appropriate. To recycle this container, please triple rinse and drop off to your local AgRecovery collection site. For details on how to correctly prepare your container and collection sites please visit the AgRecovery website [www.agrecovery.co.nz](http://www.agrecovery.co.nz)

**EMERGENCY RESPONSE** In case of spillage, wear appropriate protective clothing and prevent material from entering water ways. Absorb spills with inert material and place in waste container. Sweep up solids without creating dust. Wash area with water and absorb with further inert material. Dispose of waste safely.



**Inovata**<sup>®</sup>

Registered to: Inovata Ltd  
16 Sheffield Crescent, Burnside, Christchurch 8053  
PO Box 20449, Bishopdale, Christchurch 8543  
[www.inovata.co.nz](http://www.inovata.co.nz) | [sales@inovata.co.nz](mailto:sales@inovata.co.nz)  
Freephone 0800 942 006

