

**DANGER**  
FOR ANIMAL TREATMENT ONLY.  
KEEP OUT OF REACH OF CHILDREN.

Inovata®

# CattleAL

## DOUBLE COMBINATION ORAL DRENCH FOR CATTLE

For the control of roundworms, tapeworm, lungworm and adult liver fluke in cattle, including roundworms resistant to benzimidazole or levamisole.



MADE  
HERE.  
FOR  
HERE.



**2**

Albendazole  
Levamisole Hydrochloride

**5L**

**ACTIVE INGREDIENTS:**  
100 g/L Albendazole and 75 g/L Levamisole Hydrochloride.

**Inovata®**

# Inovata® CattleAL

## READ ENTIRE LABEL BEFORE USE

**HANDLING PRECAUTIONS** Harmful if swallowed. May cause an allergic skin reaction. Suspected of causing genetic defects. May damage fertility or the unborn child. May cause damage to organs. Wash hands thoroughly after handling. Do not eat, drink or smoke when using this product. Contaminated work clothing should not be allowed out of the workplace. Wear protective gloves/clothing. Obtain special instructions before use. Do not handle until all safety precautions have been read and understood. Use personal protective equipment as required. Do not breathe dust/fumes/gas/mist/vapours/spray.

**FIRST AID** If medical advice is needed, have product container or label at hand.

**IF SWALLOWED:** Call a POISON CENTER or doctor/physician if you feel unwell. Rinse mouth.

**IF ON SKIN:** Wash with plenty of soap and water. If skin irritation or rash occurs: Get medical advice/attention. Wash contaminated clothing before reuse. If exposed or concerned, or feel unwell: Get medical advice/attention.

For advice, contact the National Poisons Centre - 0800 POISON (0800 764 766) or a doctor immediately.

**ENVIRONMENTAL PROTECTION** Harmful to terrestrial vertebrates. Avoid release to the environment and contamination of any water supply with product or empty container. Collect spillage.

**INDICATIONS** For the control of sensitive mature and immature haemonchus, ostertagia, nematodirus, cooperia, strongyloides, bunostomum, oesophagostomum, chabertia that may be resistant to benzimidazole or levamisole. Also for the control of tapeworm, lungworm and adult liver fluke in cattle.

Resistance can develop to any anthelmintic.

Ask your local veterinary practitioner or animal health adviser for recommended parasite management practices for your area to reduce development of resistance. It is advisable that a resistance test be conducted regularly when using any parasite treatment.

**PRECAUTIONS** Doses three or more times those recommended can cause symptoms of Levamisole toxicity, so estimate liveweights carefully. Animals under stress due to advanced pregnancy, adverse weather conditions, poor nutrition, prolonged yarding or dehydration may be more susceptible to toxicity.

Fatal interactions may occur between Levamisole and organophosphate dips. Do not exceed the stated dose volume.

**DOSAGE** Cattle 1 mL per 10 kg bodyweight

Bodyweight	Dose	Doses per 5 L Pack
60 kg	6 mL	833
61-70 kg	7 mL	714
71-80 kg	8 mL	625
81-90 kg	9 mL	555
91-100 kg	10 mL	500
101-110 kg	11 mL	454
111-120 kg	12 mL	416
121-130 kg	13 mL	384
131-140 kg	14 mL	357
141-150 kg	15 mL	333
151-160 kg	16 mL	312
161-170 kg	17 mL	294
171-180 kg	18 mL	277
181-190 kg	19 mL	263
191-200 kg	20 mL	250
201-210 kg	21 mL	238
211-220 kg	22 mL	227
221-230 kg	23 mL	217
231-240 kg	24 mL	208
241-250 kg	25 mL	200
251-260 kg	26 mL	192
261-270 kg	27 mL	185
271-280 kg	28 mL	178
281-290 kg	29 mL	172
291-300 kg	30 mL	166

For animals heavier than 300 kg add 1 mL per 10 kg bodyweight.

**DIRECTIONS FOR USE** By law user must take due care, obtaining expert advice if necessary, to avoid unnecessary pain and distress when using the product other than directed on the label.

Draft animals into lines of similar bodyweight. A representative sample of animals should be weighed before treatment. Do not underdose. Where there is a large variation in size within the group, dose rate should be based on the label directions for each weight range. Drafting into two or more lines may be appropriate, to avoid excessive overdosing. Correct drenching techniques should be used. Check accuracy of drench guns regularly.

**DISPOSAL** Preferably dispose of product by use. Otherwise dispose of product and packaging at an approved landfill/facility. Triple rinse container. Recycle empty rinsed container at an approved recycling centre. Otherwise puncture or crush empty container and dispose of at an approved landfill/facility. Do not use container for any other purpose.

**STORAGE** Store locked up below 25°C. Away from sunlight. Do not freeze.

**WITHOLDING PERIODS** It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

**MILK:** Milk intended for sale for human consumption must be discarded during treatment and for not less than 35 days following the last treatment.

**MEAT:** Animals producing meat or offal for human consumption must not be sold for slaughter either during treatment or within 14 days of cessation of last treatment.

ENVIRONMENTALLY HAZARDOUS SUBSTANCE,  
LIQUID N.O.S. (Contains Albendazole), UN No.3082

Registered pursuant to the ACVM Act 1997, No A011957.  
See [www.foodsafety.govt.nz](http://www.foodsafety.govt.nz) for registration conditions.

Approved pursuant to the HSN0 Act 1996, HSRI00758.  
See [www.epa.govt.nz](http://www.epa.govt.nz) for controls.



Registered to: Inovata Ltd  
16 Sheffield Crescent, Burnside, Christchurch 8053  
PO Box 20449, Bishopdale, Christchurch 8543  
[www.inovata.co.nz](http://www.inovata.co.nz) | [sales@inovata.co.nz](mailto:sales@inovata.co.nz)  
Freephone 0800 942 006



Batch:  
Expiry: