

BEEF DRENCH PLAN

| ANIMAL HEALTH

PRODUCT	JUL	AUG	SEP	OCT	NOV	DEC	JAN	FEB	MAR	APR	MAY	JUN	
Beef Calves off Cows								Control of the major parasites of young cattle, Ostertagia, lungworm and Ivermectin resistant Cooperia					
Inovata Cattle AL								From weaning to June in fluke areas alternating with Gemini or Inovata Abamec					
Gemini								Weaning		Regular drenching every 6 weeks			
Inovata Abamectin								Lice control heading into winter					
Beef Cattle RIs	Control of the major parasites of young cattle, Ostertagia, lungworm and Ivermectin resistant Cooperia								Autumn Ostertagia, Fluke (if required) and lice control				
Gemini	Every 6 - 8 weeks												
Inovata Abamectin								Autumn treatment when larval challenge is high and lice control					
PRODUCT	JUL	AUG	SEP	OCT	NOV	DEC	JAN	FEB	MAR	APR	MAY	JUN	
Dairy Calves & Weaners				Control of the major parasites of young cattle, Ostertagia, lungworm and Ivermectin resistant Cooperia									
Inovata Cattle AL				Weaning				Use in fluke areas alternating with Gemini					
Gemini				Post Weaning every 4 - 6 weeks Calves > 100kg									
Inovata Abamectin													
Dairy Heifers	Control of the major parasites of young cattle, Ostertagia, lungworm and Ivermectin resistant Cooperia				Autumn Ostertagia, Fluke (if required) and lice control								
Gemini	Every 6 - 8 weeks												
Inovata Abamectin								Autumn treatment when larval challenge is high and lice control					
Lactating Dairy Cows													
Inovata Abamectin											Cost effective lice control at dry off		

DAIRY DRENCH PLAN

ANIMAL HEALTH

PRODUCT	JUL	AUG	SEP	OCT	NOV	DEC	JAN	FEB	MAR	APR	MAY	JUN
Dairy Calves & Weaners	Control of the major parasites of young cattle, Ostertagia, lungworm and Ivermectin resistant Cooperia											
Inovata Cattle AL				Weaning					Use in fluke areas alternating with Gemini			
Gemini				Post Weaning every 4 - 6 weeks Calves > 100kg								
Inovata Abamectin												
Dairy Heifers	Control of the major parasites of young cattle, Ostertagia, lungworm and Ivermectin resistant Cooperia					Autumn Ostertagia, Fluke (if required) and lice control						
Gemini	Every 6 - 8 weeks											
Inovata Abamectin									Autumn treatment when larval challenge is high and lice control			
Lactating Dairy Cows												
Gemini												
Inovata Abamectin										Cost effective lice control at dry off		

SHEEP DRENCH PLAN | ANIMAL HEALTH

PRODUCT	JUL	AUG	SEP	OCT	NOV	DEC	JAN	FEB	MAR	APR	MAY	JUN			
Lambs	Consider exit drenching		Use a pre weaning drench to increase growth rates pre weaning.				Drench every 28 days. Consider Haemonchus control in effected areas.				Monitor drench efficacy against small intestinal Trichostrongylus				
Inovata Tape				Pre weaning Coordinate with your 5 n 1 vaccination policy		Weaning									
Eraser Tape															
Inovata OXL	Late Autumn and Winter									Post weaning				Late Autumn and Winter	
Gemini															
Eraser															
Eraser OX															
Tison															
Adult Sheep	Ewes 2 - 3 weeks prior to lambing										Ewes 2 - 3 weeks prior to lambing				
Inovata Tape															
Eraser Tape															
Inovata OXL	Pre Lamb Tailor drench choice to ewe drenching requirements and the drench resistance status of the farm		Docking					Consider the requirement for Haemonchus control in younger and lighter condition ewes			Pre Lamb				
Gemini															
Eraser															
Eraser OX															
Tison															
Fly Strike Control in Sheep		Spray On = Lambs at docking/tailing, convenient use Liquid = When flystrike is eminent, for use with a jetting race or shower dip													
Cyro-Spray		Lambs at docking/tailing													
Cyro-Liquid					Warm and humid conditions, flystrike prevention.										