

WARNING

READ LABEL BEFORE USE.
KEEP OUT OF REACH OF CHILDREN.
FOR ANIMAL TREATMENT ONLY.


Inovata®

Selenium Pour-on

For the prevention and treatment of
selenium deficiency and **selenium**
responsive diseases in **cattle** and **deer**.

1L

MADE
HERE.
FOR
HERE.



ACTIVE INGREDIENT:

5 mg/mL **Selenium** (as Sodium Selenate).
Contains: 300 g/L Diethylene Glycol Monobutyl Ether.

Inovata®

READ THE ENTIRE LABEL BEFORE USE

Read label and obtain special instructions before use. Do not handle until all safety precautions have been read and understood.

HANDLING PRECAUTIONS: Harmful if inhaled. Causes serious eye irritation. Suspected of causing genetic defects. May cause damage to organs through prolonged or repeated exposure.

SAFETY PRECAUTIONS AND DIRECTIONS: Do not handle until all safety precautions have been read and understood. Wash hands and exposed skin thoroughly after handling, before meals and after use. Wear protective gloves/protective clothing/eye protection. Do not breathe mists/fumes/spray/vapours. Use only outdoors or in a well ventilated area.

FIRST AID: If inhaled, remove person to fresh air and keep comfortable for breathing. For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If in eyes, rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists, get medical advice/attention. If exposed, you feel unwell or concerned, get medical advice/attention.

ENVIRONMENTAL PROTECTION: Very toxic to aquatic life with long lasting effects. Avoid release to the environment. Collect spillage.

EMERGENCY RESPONSE: In case of spillage, wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Wash the area with water and absorb with further inert material. Dispose of waste safely.

CONTRAINDICATIONS: Do not use on cattle with light coats intended for showing. A brown discolouration can occur. Do not use where **selenium** deficiency has not been diagnosed. Do not use at the same time as any other selenised fertiliser, prill or product without consulting a veterinarian. Do not exceed stated dose, or dose more frequently than 3-weekly.

SYMPTOMS OF POISONING: Nausea, garlic odour of breath, nervousness, gastrointestinal upset.

DISPOSAL: Preferably dispose of the product by use. Otherwise dispose of product and packaging at an approved landfill or other approved facility. Avoid contamination of any water source. Triple rinse container immediately after emptying and pour rinsate onto waste ground. Recycle via AGRECOVERY Rural Recycling Programme (see www.agrecovery.co.nz). Do NOT use container for any other purpose.

STORAGE: Store locked up below 25°C. Must be discarded 56 days (8 weeks) after first use.

WITHHOLDING PERIODS: It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

MILK: Nil **MEAT:** Nil

Inovata®

Selenium Pour-on

DIRECTIONS FOR USE: By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

DOSAGE AND ADMINISTRATION: SHAKE THOROUGHLY BEFORE USE.

Apply along the backline from between the shoulders to the tail.
Not to be administered orally.

CATTLE: 1.5 mL/50 kg bodyweight (0.15 mg Se per kg)

DEER: 3 mL/50 kg bodyweight (0.3 mg Se per kg)

CONTRAINDICATIONS:

Do not use on cattle with light coats intended for showing. A brown discolouration can occur. Do not use where selenium deficiency has not been diagnosed.

Do not use at the same time as any other selenised fertiliser, prill or product without consulting a veterinarian.
Do not exceed stated dose, or dose more frequently than 3-weekly.

SYMPTOMS OF POISONING: Nausea, garlic odour of breath, nervousness, gastrointestinal upset.

DISPOSAL: Preferably dispose of the product by use. Otherwise dispose of product and packaging at an approved landfill or other approved facility. Avoid contamination of any water source. Triple rinse container immediately after emptying and pour rinsate onto waste ground. Recycle via AGRECOVERY Rural Recycling Programme (see www.agrecovery.co.nz). Do NOT use container for any other purpose.

STORAGE: Store locked up below 25°C. Must be discarded 56 days (8 weeks) after first use.

WITHHOLDING PERIODS: It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice:
Maximum Residue Levels for Agricultural Compounds.

MILK: Nil

MEAT: Nil



MORE INFO



| CATTLE | | |
|---------------------------|---------|------------|
| Body-weight | Dose | 1 L treats |
| 50 kg | 1.5 mL | 666 |
| 100 kg | 3 mL | 333 |
| 150 kg | 4.5 mL | 222 |
| 200 kg | 6 mL | 166 |
| 250 kg | 7.5 mL | 133 |
| 300 kg | 9 mL | 111 |
| 350 kg | 10.5 mL | 95 |
| 400 kg | 12 mL | 83 |
| 450 kg | 13.5 mL | 74 |
| 500 kg | 15 mL | 66 |
| Above 500 kg 1.5 mL/50 kg | | |

| DEER | | |
|-------------------------|-------|------------|
| Body-weight | Dose | 1 L treats |
| 50 kg | 3 mL | 333 |
| 100 kg | 6 mL | 166 |
| 150 kg | 9 mL | 111 |
| 200 kg | 12 mL | 83 |
| 250 kg | 15 mL | 66 |
| 300 kg | 18 mL | 55 |
| 350 kg | 21 mL | 47 |
| 400 kg | 24 mL | 41 |
| 450 kg | 27 mL | 37 |
| 500 kg | 30 mL | 33 |
| Above 500 kg 3 mL/50 kg | | |

CONTAINS: Sodium Selenate 12.4 g/L, Diethylene Glycol Monobutyl Ether 300 g/L

Registered to: Inovata Ltd

16 Sheffield Crescent, Burnside, Christchurch 8053
www.inovata.co.nz | sales@inovata.co.nz | Freephone: 0800 942 006

Registered pursuant to the ACVM Act 1997: A012120
See www.foodsafety.govt.nz for registration conditions.

PEEL HERE FOR FURTHER INFORMATION

ZPLAB528 VER-1

Inovata[®]

Selenium Pour-on

DIRECTIONS FOR USE: By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

DOSAGE AND ADMINISTRATION: SHAKE THOROUGHLY BEFORE USE.

Apply along the backline from between the shoulders to the tail.
Not to be administered orally.

CATTLE: 1.5 mL/50 kg bodyweight (0.15 mg Se per kg)

DEER: 3 mL/50 kg bodyweight (0.3 mg Se per kg)

| CATTLE | | |
|---------------------------|---------|------------|
| Body-weight | Dose | 1 L treats |
| 50 kg | 1.5 mL | 666 |
| 100 kg | 3 mL | 333 |
| 150 kg | 4.5 mL | 222 |
| 200 kg | 6 mL | 166 |
| 250 kg | 7.5 mL | 133 |
| 300 kg | 9 mL | 111 |
| 350 kg | 10.5 mL | 95 |
| 400 kg | 12 mL | 83 |
| 450 kg | 13.5 mL | 74 |
| 500 kg | 15 mL | 66 |
| Above 500 kg 1.5 mL/50 kg | | |

| DEER | | |
|-------------------------|-------|------------|
| Body-weight | Dose | 1 L treats |
| 50 kg | 3 mL | 333 |
| 100 kg | 6 mL | 166 |
| 150 kg | 9 mL | 111 |
| 200 kg | 12 mL | 83 |
| 250 kg | 15 mL | 66 |
| 300 kg | 18 mL | 55 |
| 350 kg | 21 mL | 47 |
| 400 kg | 24 mL | 41 |
| 450 kg | 27 mL | 37 |
| 500 kg | 30 mL | 33 |
| Above 500 kg 3 mL/50 kg | | |

PHIL STC MULTIPLE[™]



CONTAINS: Sodium Selenate 12.4 g/L, Diethylene Glycol Monobutyl Ether 300 g/L

Registered pursuant to the
ACVM Act 1997: A012120
See www.foodsafety.govt.nz
for registration conditions.

