WARNING

READ LABEL BEFORE USE.
KEEP OUT OF REACH OF CHILDREN.
FOR ANIMAL TREATMENT ONLY.

Inovata

Selenium Pour-on

For the prevention and treatment of **selenium** deficiency and **selenium** responsive diseases in **cattle** and **deer**.

2.5L

ACTIVE INGREDIENT: 5 mg/mL **Selenium** (as Sodium Selenate). Contains: 300 g/L Diethylene Glycol Monobutyl Ether.

Inovata[®]

Selenium Pour-on

READ THE ENTIRE LABEL BEFORE USE

Read label and obtain special instructions before use. Do not handle until all safety precautions have been read and understood.

HANDLING PRECAUTIONS: Harmful if inhaled. Causes serious eye irritation. Suspected of causing genetic defects. May cause damage to organs through prolonged or repeated exposure.

SAFETY PRECAUTIONS AND DIRECTIONS: Do not handle until all safety precautions have been read and understood. Wash hands and exposed skin thoroughly after handling, before meals and after use. Wear protective gloves/protective clothing/eye protection. Do not breathe mists/fumes/spray/vapours. Use only outdoors or in a well ventilated area.

FIRST AID: If inhaled, remove person to fresh air and keep comfortable for breathing. For advice, contact the National Poisons Centre 0800 POISON (0800 764 766) or a doctor immediately. If in eyes, rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists, get medical advice/attention. If exposed, you feel unwell or concerned, get medical advice/attention.

ENVIRONMENTAL PROTECTION: Very toxic to aquatic life with long lasting effects. Avoid release to the environment. Collect spillage.

EMERGENCY RESPONSE: In case of spillage, wear appropriate protective clothing and prevent material from entering waterways. Absorb spills with inert material and place in waste containers. Wash the area with water and absorb with further inert material. Dispose of waste safely.

DIRECTIONS FOR USE: By law the user must take due care, obtaining expert advice when necessary, to avoid unnecessary pain and distress when using the product other than as directed on the label.

DOSAGE AND ADMINISTRATION: SHAKE THOROUGHLY BEFORE USE.

Apply along the backline from between the shoulders to the tail. Not to be administered orally.

CATTLE: 1.5 mL/50 kg bodyweight (0.15 mg Se per kg) **DEER:** 3 mL/50 kg bodyweight (0.3 mg Se per kg)

CATTLE			
Body- weight	Dose	2.5 L treats	
50 kg	1.5 mL	1666	
100 kg	3 mL	833	
150 kg	4.5 mL	555	
200 kg	6 mL	416	
250 kg	7.5 mL	333	
300 kg	9 mL	277	
350 kg	10.5 mL	238	
400 kg	12 mL	208	
450 kg	13.5 mL	185	
500 kg	15 mL	166	
Above 5	00 kg 1.5 m	L/50 kg	

DEER		
Body- weight	Dose	2.5 L treats
50 kg	3 mL	833
100 kg	6 mL	416
150 kg	9 mL	277
200 kg	12 mL	208
250 kg	15 mL	166
300 kg	18 mL	138
350 kg	21 mL	119
400 kg	24 mL	104
450 kg	27 mL	92
500 kg	30 mL	83
Above 500 kg 3 mL/50 kg		

CONTAINS: Sodium Selenate 12.4 g/L, Diethylene Glycol Monobutyl Ether 300 g/L

Registered to:

Inovata Ltd, 16 Sheffield Crescent, Burnside, Christchurch 8053 www.inovata.co.nz | sales@inovata.co.nz | Freephone 0800 942 006 **CONTRAINDICATIONS:** Do not use on cattle with light coats intended for showing. A brown discolouration can occur. Do not use where **selenium** deficiency has not been diagnosed. Do not use at the same time as any other selenised fertiliser, prill or product without consulting a veterinarian. Do not exceed stated dose, or dose more frequently than 3-weekly.

SYMPTOMS OF POISONING: Nausea, garlic odour of breath, nervousness, gastrointestinal upset.

DISPOSAL: Preferably dispose of the product by use. Otherwise dispose of product and packaging at an approved landfill or other approved facility. Avoid contamination of any water source. Triple rinse container immediately after emptying and pour rinsate onto waste ground. Recycle via AGRECOVERY Rural Recycling Programme (see www.agrecovery.co.nz). Do NOT use container for any other purpose.

STORAGE: Store locked up below 25°C. Must be discarded 56 days (8 weeks) after first use.

WITHHOLDING PERIODS: It is an offence for users of this product to cause residues exceeding the relevant MRL in the Food Notice: Maximum Residue Levels for Agricultural Compounds.

MILK: Nil MEAT: Nil

Registered pursuant to the ACVM Act 1997: A012120 See www.foodsafety.govt.nz for registration conditions.









Batch: Expiry:

ZPLAB529 VER-1